

Deaths and Case Fatality Ratios for Select Vaccine Preventable Diseases by Report Year, Colorado, 2000 – 2009*

YEAR	Invasive <i>Haemophilus influenzae</i> Serotype b (Hib)			Invasive Meningococcal Disease			Pertussis			Invasive Pneumococcal Disease [§]		
	Total	Deaths	CFR ^{§§}	Total	Deaths	CFR ^{§§}	Total	Deaths	CFR ^{§§}	Total	Deaths	CFR ^{§§}
2000	3	1	33.3	35	4	11.4	487	2	0.4			
2001	4	1	25.0	37	4	10.8	389	0	0.0	317	45	14.2
2002	2	1	50.0	24	1	4.2	464	0	0.0	390	46	11.8
2003	0	0	0.0	27	5	18.5	368	0	0.0	367	46	12.5
2004	1	0	0.0	15	2	13.3	1185	1	0.1	331	40	12.1
2005	2	0	0.0	19	3	15.8	1383	2	0.1	420	45	10.7
2006	0	0	0.0	22	1	4.5	710	0	0.0	419	39	9.3
2007	2	0	0.0	22	2	9.0	306	0	0.0	458	40	8.7
2008	0	0	0.0	16	2	12.5	160	0	0.0	541	32	5.9
2009 ⁺⁺	3	0	0.0	24	3	12.5	201	0	0.0	623	52	8.3
Annual Average	1.7	0.3	17.6	24.1	2.7	11.2	565.3	0.5	0.1	443.6	42.5	9.6

*There were no diphtheria, measles, mumps, rubella, or tetanus associated deaths reported January 1, 2000 – December 31, 2009

[§]Invasive pneumococcal disease became reportable statewide December 1, 2001. Annual average is a 8-year average (2002 – 2009).

^{§§}CFR=Case Fatality Ratio (expressed as a percent) [(#Deaths/#Cases)*100].

⁺In 2001 one infant died of co-infection with *Haemophilus influenzae* and pneumococcal disease. This infant is counted in statistics for both diseases.

⁺⁺ 2009 death information is provisional for invasive *Haemophilus influenzae*, meningococcal and pneumococcal diseases.