

















2009 H1N1 Influenza Vaccine Dosage Chart



PLEASE TRIPLE CHECK ALL VACCINATIONS BEFORE ADMINISTERING



Manufacturer	Age	Dose-Presentation	Number of Doses Interval	Route
	Children (4 through 9 years)	0.5mL prefilled syringe or multidose vial	2* with 28 day interval	Intramuscular 
	Children (10 through 17 years)	0.5mL prefilled syringe or multidose vial	1	Intramuscular 
	Adults (18 years & older)	0.5mL prefilled syringe or multidose vial	1	Intramuscular 
	Children (6 through 35 months)	0.25mL prefilled syringe or multidose vial	2* with 28 day interval	Intramuscular 
	Children (3 through 9 years)	0.5mL prefilled syringe or multidose vial	2* with 28 day interval	Intramuscular 
	Children (10 years & older)	0.5mL prefilled syringe or multidose vial	1	Intramuscular 
	Adults	0.5mL prefilled syringe or multidose vial	1	Intramuscular 
	3 years of age & older ▲	0.5mL prefilled syringe or multidose vial	1*	Intramuscular 
	Adults 18 years of age & older	0.5 mL prefilled syringe or multidose vial	1	Intramuscular 
	Children (2 through 9 years)	0.2 mL Spray 0.1mL (1/2 dose) into each nostril as indicated on the syringe.	2*◇ with 28 day interval	Intranasal 
	Children, adolescents and adults (10 through 49 years)	0.2mL Spray 0.1mL (1/2 dose) into each nostril as indicated on the syringe.	1◇	Intranasal 

★If different types of vaccine are used for the first and booster dose, there is no need to revaccinate a child.

◇Two doses of LAIV (Seasonal and 2009H1N1) should not be co-administered and should be separated by at least one month (28 days).

▲ CSL License status changed 11/12/09 now licensed for ≥age 6 months BUT is recommended for use in children/persons ≥ 3 years. If used for children ages 6-35 months the dose must be decreased to 0.25mL