

## Bonny Reservoir – Summary of Hg Concentrations

118 fish sampled = 21 Common Carp  
2 Wiper  
18 Freshwater Drum  
51 Walleye  
3 Black Crappie  
2 White Bass  
15 Gizzard Shad  
6 White Sucker

34 Composite Samples = 4 Common Carp, composited 3 fish per sample  
2 Wiper, composited 1 fish per sample  
6 Freshwater Drum, composited 3 fish per sample  
12 Walleye, composited 5 fish per sample  
1 Black Crappie, composited 3 fish per sample  
2 White Bass, composited 1 fish per sample  
5 Gizzard Shad, composited 3 fish per sample  
2 White Sucker, composited 3 fish per sample

### Results:

For the Common Carp samples = 0.13, 0.21, 0.13, 0.12 = all below 0.5

For the Wiper samples = 0.27, 0.13 = all below 0.5

For the Freshwater Drum samples = <0.1, <0.1, <0.1, <0.1, 0.13, 0.18 = all below 0.5

For the Walleye samples = <0.1, <0.1, <0.1, <0.1, <0.1, <0.1, <0.1, <0.1, <0.1, <0.1, <0.1, 0.11 = all below 0.5

For the Black Crappie samples = <0.1 = all below 0.5

For the White Bass samples = <0.1, <0.1 = all below 0.5

For the Gizzard Shad samples = <0.1, <0.1, <0.1, <0.1, <0.1 = all below 0.5

For the White Sucker samples = <0.1, <0.1 = all below 0.5